



MARIANNE MORELLI, D.M.D.

BROOKFIELD FAMILY DENTISTRY, LLC

## Post-Operative Instructions for Gingival Grafting

**Bleeding** The wound may ooze for up to 24 hours. If oozing continues for more than 24 hours please call the office during normal business hours or the doctor directly if the office is closed.

**Ice** Ice will help with bleeding, swelling and pain. Swelling is a normal reaction of the irritated tissue. Apply cold packs to the face adjacent to surgical area as soon as possible for the first 6-12 hours post-operatively. Ten minutes on and ten minutes off is recommended. Usually swelling is more pronounced during the first 48 hours.

**Rest** Go home and relax for the remainder of the day. Strenuous activity and exercise should be avoided for 72 hours. Routine activity is fine beginning the second day.

**Home Care** Do not brush directly on the wound for the two weeks. **Leave it alone!** Do not rinse or spit – let any saliva gently dribble out. It is normal to have a funny taste and/or smell the two weeks. You may gently rinse with warm salt water. In addition you may be directed to gently rinse with a prescription mouth wash. The use of C-PAP or Night Guard appliances should be continued as directed, unless otherwise noted.

**Diet** For a period of two weeks avoid hard, crunchy, chewy and particulate foods. Try to eat soft foods such as eggs, pasta, and yogurt and take small mouthfuls chewed carefully and slowly on the opposite side. Avoid very hot foods and beverages, spicy foods, sharp foods and salty foods. No drinking through a straw for two weeks.

**Medications** If a medication has been prescribed take all exactly as prescribed and complete the course of treatment.

**Antibiotic:** Antibiotics are used for specific procedures and for individuals with specific requirements. Take them all and exactly as prescribed.

**Pain:** Most often anti-inflammatory agents NSAIDS (Advil/Motrin) and/or acetaminophen (Tylenol) will be enough to control your discomfort. If necessary an additional prescription will be given to alleviate pain. Avoid alcohol consumption with narcotics.

**Smoking and Alcohol** Avoid both for as long as possible. It is well documented both cause a significant increase in post-operative complications such as infection, bleeding, delayed healing and poor outcome.

### **OTC PAIN MEDICATION REGIMEN:**

**Take 3 Advil (200 mg Ibuprofen) taken with 2 Regular Strength Tylenol all at the same time, taken every 6 hours for up to 72 hours**

**Special Instructions:** \_\_\_\_\_

**Call us promptly in case of severe pain, excessive bleeding or any unusual developments.**

**Office: 203-775-6167 Dr. Morelli: 203-448-6469**

DIPLOMATE,  
AMERICAN BOARD OF PERIODONTOLOGY  
AND IMPLANT DENTISTRY



834 FEDERAL ROAD, BROOKFIELD, CT 06804  
EMAIL: OFFICE@DRMORELLI.COM  
OFFICE: (203)775-6167  
FAX: (203)775-6169